

## On the role of v-ATPase V0a1—dependent degradation in Alzheimer disease

W. Ryan Williamson and P. Robin Hiesinger\*

Department of Physiology and Green Center for Systems Biology; University of Texas Southwestern Medical Center at Dallas; Dallas, TX USA

Defective autophagy and lysosomal degradation are hallmarks of numerous neurodegenerative disorders. Vesicular ATPases are intracellular proton pumps that acidify autophagosomes and lysosomes. V0a1 is a key component of the v-ATPase that is only required in neurons in *Drosophila melanogaster*. We have recently shown that loss of V0a1 in *Drosophila* photoreceptor neurons leads to slow, adult-onset degeneration.<sup>1</sup> Concurrently, Lee et al.<sup>2</sup> reported that V0a1 fails to localize to lysosomal compartments in cells from Presenilin 1 knock-out cells. Together these two reports suggest that a neuronal V0a1-dependent degradation mechanism may be causally linked to Alzheimer pathology. Indeed, we now show that loss of V0a1 makes *Drosophila* neurons more susceptible to insult with human Alzheimer-related neurotoxic A $\beta$  and tau proteins. Furthermore, we discuss the potential significance of the discovery of the neuron-specific degradation mechanism in *Drosophila* for intracellular degradation defects in Alzheimer Disease.

increase of undegraded endosomal and autophagosomal compartments, and ultimately to slow, adult-onset degeneration.<sup>1</sup> Furthermore, loss of *v0a1* also causes severe neurotransmission defects and thereby embryonic lethality. The recent *Drosophila* studies were made possible by a genetic method that only renders photoreceptor neurons homozygous mutant in otherwise heterozygous animals.<sup>3</sup> Since photoreceptors are not required for viability, this system allows one to overcome early neuronal death due to embryonic lethality and follow the maintenance and degeneration of homozygous mutant neurons in an otherwise healthy fly. *v0a1* photoreceptors exhibit loss of neurotransmission immediately after eclosion,<sup>6</sup> whereas significant functional and morphological neurodegeneration occurs only after 1–2 weeks.<sup>1</sup> Importantly, *v0a1* is only required in neurons in *Drosophila*.<sup>6</sup> The discovery of a neuron-specific degradation mechanism in *Drosophila* establishes a causal link between intracellular degradation and adult-onset degeneration. Similarly, disruption of basal autophagy in mice through either loss of *atg5* or *atg7* leads to neurodegeneration within weeks,<sup>7,8</sup> and induction of autophagy can alleviate neurodegenerative effects.<sup>9,10</sup> Lysosomal storage disorders show how mutants in endolysosomal and autophagic pathways lead to late onset neurodegeneration.<sup>11</sup> Finally, neurons in mouse models of Down Syndrome and Alzheimer Disease exhibit early accumulations of Rab5-marked early endosomes as well as late endosomes.<sup>3</sup> All these findings are consistent with the idea that loss of degradation capacity sensitizes neurons

**Key words:** AUTHOR: Please provide 5–7 key words

Submitted: 08/19/10

Accepted: 08/19/10

Previously published online:

[www.landesbioscience.com/journals/cib/article/13364](http://www.landesbioscience.com/journals/cib/article/13364)

\*Correspondence to: P. Robin Hiesinger;  
Email: [robin.hiesinger@utsouthwestern.edu](mailto:robin.hiesinger@utsouthwestern.edu)

Addendum to: Williamson WR, Wang D, Haberman AS, Hiesinger PR. A dual function of V0-ATPase a1 provides an endolysosomal degradation mechanism in *Drosophila melanogaster* photoreceptors. *J Cell Biol* 2010; 189:885–99; PMID: 20513768; DOI: 10.1083/jcb.201003062.

Until recently, the degradation machinery implicated in neuronal degeneration was thought to be ubiquitous and there was little evidence for a dedicated neuronal pathway. In addition, the causal relationship between adult-onset neurodegeneration and accumulations of undegraded proteins in Alzheimer Disease as well as other neurodegenerative disorders is unclear.<sup>3,4</sup> We have recently shown that loss of the v-ATPase component V0a1 leads to a ~50% reduction of strongly acidified compartments in neurons, an

to ‘cargo overload’ as a cause for slow, adult-onset degeneration.

Lee et al.<sup>2</sup> recently reported that mislocalization of V0a1 is at least partially responsible for causing degenerative phenotypes in Presenilin 1 (PS1) knock-out cells. This finding is consistent with our discovery of the V0a1-dependent neuronal degradation mechanism in *Drosophila* and the degenerative phenotypes caused by loss of *v0a1*. The *Drosophila*, mouse and human V0a1 genes are precise orthologs within the V0a gene families in each of these different species that comprise exactly four V0a1-4 genes. Since loss of *v0a1* in *Drosophila* causes embryonic lethality due to neurotransmission failure, and neurotransmission has been shown to be preserved in PS1 KO neurons,<sup>12</sup> we conclude that the trafficking defect of V0a1 reported in PS1 KO cells should not represent a full *v0a1* loss-of-function. Instead, we propose that in PS1 KO cells only the trafficking of a subset of degradative compartments is affected. Furthermore, since lethality due to loss of neurotransmission precedes and masks slow degeneration, it may be unlikely to find disease-related *v0a1* mutations in humans. In contrast, loss of Presenilin 1 may cause a restricted loss of V0a1 in a specialized degradation pathway.

We were intrigued by the possibility that loss of *v0a1*-dependent degradation renders neurons more susceptible to neurotoxic insults. To directly test the idea that neurons lacking the neuron-specific degradation pathway are more susceptible to neurotoxic proteins, we designed challenge experiments with human A $\beta$ <sup>33</sup> and tau<sup>14</sup> variants. A $\beta$  peptides are the neurotoxic cleavage products of Amyloid Precursor Proteins (APP) that are causally associated with Alzheimer Disease and may have neurotoxic effects by disrupting lysosomal function.<sup>15</sup> Of the two possible A $\beta$  cleavage products of APP, A $\beta$ 42 is more toxic and its expression in *Drosophila* photoreceptors leads to early cell death, whereas expression of A $\beta$ 40 has much milder phenotypes.<sup>13</sup> We established a protocol using A $\beta$ 40 expression and *v0a1* RNAi expression in addition to two days intense light stimulation; under these conditions, neither A $\beta$ 40 nor *v100* RNAi cause significant degeneration as


measured by ERG response amplitudes (Fig. 1A and B). Knock-down of *v0a1* using RNAi causes the same phenotypes as loss of *v100* in photoreceptors (Fig. 2), and it provides a sensitized system for degradative challenges, as shown below. Indeed, when we co-express *v0a1* RNAi and A $\beta$ 40 under these conditions, ERG response amplitudes are significantly reduced compared to both A $\beta$ 40 or *v100* RNAi alone (Fig. 1A and B). Next, we expressed the human microtubule-binding protein tau as well as a mutant version of tau (R406W) that has been implicated in the pathogenesis of Alzheimer’s disease and related disorders.<sup>14</sup> Similar to A $\beta$ 40, expression of tau and tau<sup>R406W</sup> lead to increased functional degeneration when co-expressed with *v0a1* RNAi (Fig. 1A and B). Morphologically, two day intense light stimulation already leads to a few minor disruptions of the photoreceptor arrangement in the eye that are apparent as ‘holes’, i.e., unlabeled spaces between the rhabdomeric structures (Fig. 1C). These aberrations are increased by *v100* RNAi expression and indicative of early stages of photoreceptor degeneration. Co-expression of either A $\beta$ 40 or tau leads to a dramatic acceleration of this degenerative phenotype (Fig. 1C). These results support the idea that loss of *v0a1*-dependent degradation increases neuronal sensitivity to neurotoxic insults and accelerates neurodegeneration. Taken together with the mouse studies, loss of Presenilin 1 may cause a double burden by decreasing the degradative capacity through mislocalization of V0a1 and increasing the degradative load through faulty APP cleavage, A $\beta$  accumulations or tau tangles.

Does V0a1 define a neuron-specific degradation pathway? In *Drosophila*, V0a1 is only required in neurons which lead to the idea of a neuron-specific mechanism.<sup>1</sup> In contrast, Lee et al.<sup>2</sup> characterize a V0a1-dependent acidification defect in PS1 KO mouse blastocysts. Degradative compartments require acidification for the maturation of Cathepsin proteases. PS1 KO blastocysts have a defect in Cathepsin D maturation, although a similar defect has not yet been demonstrated for neurons.<sup>2,16</sup> Furthermore, Presenilin double knock-out hippocampal neurons have significant ER

calcium signaling abnormalities which may also potentially be linked to defects in V0a1 trafficking and impaired acidification of intracellular compartments.<sup>17</sup> Overall, the data from Lee et al.<sup>2</sup> suggest that a degradation mechanism that is neuron-specific in *Drosophila* may also operate in non-neuronal cells in mammals. Indeed, expression data inferred from Expressed Sequence Tag sources available from the public database of the National Center for Biotechnology Information (NCBI) indicates that *v0a1* expression is not restricted to neurons in mouse and human. Interestingly, the observation of neuronal specificity for a protein that is more widely expressed in mammals has also been made for APPL, the *Drosophila* ortholog of the A $\beta$ -precursor protein APP. While expression of *Drosophila* APPL is restricted to the nervous system,<sup>18,19</sup> two of the three APP homologs in the mouse are ubiquitously expressed.<sup>20</sup> However, the recent findings in *v0a1* *Drosophila* neurons and PS1 mouse KO cells suggest that *v0a1* is not required for all intracellular degradation. Instead, V0a1 increases degradative capacity and its loss alone is sufficient to cause adult-onset degeneration in *Drosophila* neurons. In addition, loss of V0a1-dependent degradation makes cells more vulnerable to neurotoxic insults including A $\beta$  and tau proteins. Further experiments will be needed to validate the contribution of the *v0a1*-dependent degradation mechanism to neurodegenerative pathology.

## Materials and Methods

Experiments were conducted using flies of the genotype UAS-Dicer2;GMR-Gal4;UAS-*v0a1*-RNAi and additionally expressing either UAS-A $\beta$ 40, UAS-tau or UAS-tau<sup>406</sup>. The *v0a1* RNAi lines was obtained from the VDRC.<sup>21</sup> Electroretinogram recordings, immunohistochemistry and quantification were all conducted as described in Williamson et al.<sup>1</sup> Dissections were performed as described in Williamson and Hiesinger.<sup>22</sup> For immunohistochemistry, the following antibodies and dilutions were used: Choptin (mAb 24B10) at 1:50, Avalanche/Syx7 at 1:1,000 and Sunglasses at 1:1,000.




**Figure 1.** Loss of the *v0a1*-dependent degradation pathway increases susceptibility to neurotoxic insults by human A $\beta$  and tau proteins. (A) Representative ERG recording for at least  $n = 20$  three day-old flies of the genotypes indicated and raised at room temperature under constant light stimulation. (B) Quantification of ERG depolarizations. Orange bar shows ERG depolarization for control or overexpressed proteins in wild type photoreceptors; blue bars are corresponding experiments with dicer2-enhanced *v0a1* RNAi. Asterisks show statistical significance in pairwise comparisons ( $p < 0.005$ ). (C) Immunolabeled adult eyes from experimental flies used for ERG recordings shown in (A). Chaoptin and Toto-3 labeling as in Figure; Syx7/Avl labeling in red. Arrows indicate degenerative 'holes' in the immunolabeling. Arrowheads indicate early endosomal Syx7-positive accumulations with Chaoptin that are typical for loss of *v0a1* function. Error bars are S.E.M.

#### Acknowledgements

We would like to thank Drs. Helmut Kramer, Damian Crowther, Craig Montell, the Bloomington Stock Center, and the University of Iowa Developmental Studies Hybridoma Bank for reagents. We further thank Drs. Ilya Bezprozvanny and Adam Haberman for comments on the manuscript and discussion. This work was supported by grants from the National Institute of Health (RO1EY018884) and the Welch Foundation (I-1657). P.R.H. is a Eugene McDermott Scholar in Biomedical Research.

#### References

- Williamson WR, Wang D, Haberman AS, Hiesinger PR. A dual function of V0-ATPase a1 provides an endolysosomal degradation mechanism in *Drosophila* photoreceptors. *J Cell Biol* 2010.
- Lee JH, Yu WH, Kumar A, Lee S, Mohan PS, Peterhoff CM, et al. Lysosomal proteolysis and autophagy require presenilin 1 and are disrupted by Alzheimer-related PS1 mutations. *Cell* 2010; 141:1146-58.
- Nixon RA, Yang DS, Lee JH. Neurodegenerative lysosomal disorders: a continuum from development to late age. *Autophagy* 2008; 4:590-9.
- Wong E, Cuervo AM. Autophagy gone awry in neurodegenerative diseases. *Nat Neurosci* 2010; 13:805-11.
- Newsome TP, Asling B, Dickson BJ. Analysis of *Drosophila* photoreceptor axon guidance in eye-specific mosaics. *Development* 2000; 127:851-60.
- Hiesinger PR, Fayyazuddin A, Mehta SQ, Rosenmund T, Schulze KL, Zhai RG, et al. The v-ATPase V0 subunit a1 is required for a late step in synaptic vesicle exocytosis in *Drosophila*. *Cell* 2005; 121:607-20.
- Komatsu M, Waguri S, Chiba T, Murata S, Iwata J, Tanida I, et al. Loss of autophagy in the central nervous system causes neurodegeneration in mice. *Nature* 2006; 441:880-4.
- Hara T, Nakamura K, Matsui M, Yamamoto A, Nakahara Y, Suzuki-Migishima R, et al. Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice. *Nature* 2006; 441:885-9.
- Wang T, Lao U, Edgar BA. TOR-mediated autophagy regulates cell death in *Drosophila* neurodegenerative disease. *J Cell Biol* 2009; 186:703-11.
- Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature* 2008; 451:1069-75.



**Figure 2.** *v0a1* RNAi in photoreceptors causes phenotypes identical to loss of *v0a1*. (A–F) Antibody-labeling of Syx7/Avl and Sunglasses reveals similar accumulations of both markers in photoreceptors expressing dicer2-enhanced *v0a1* RNAi (C and D) as in *v0a1* null mutant photoreceptors in *eyFLP v0a1* flies (E and F). (G) Both *v0a1* RNAi as well as *ey35FLP v0a1* lead to a complete loss of neurotransmission, as indicated by the loss of electroretinogram (ERG) 'On' transients.

- Futerman AH, van Meer G. The cell biology of lysosomal storage disorders. *Nat Rev Mol Cell Biol* 2004; 5:554–65.
- Feng R, Rampon C, Tang YP, Shrom D, Jin J, Kyin M, et al. Deficient neurogenesis in forebrain-specific presenilin-1 knockout mice is associated with reduced clearance of hippocampal memory traces. *Neuron* 2001; 32:911–26.
- Crowther DC, Kinghorn KJ, Miranda E, Page R, Curry JA, Duthie FA, et al. Intraneuronal Aβeta, non-amyloid aggregates and neurodegeneration in a *Drosophila* model of Alzheimer's disease. *Neuroscience* 2005; 132:123–35.
- Wittmann CW, Wszolek MF, Shulman JM, Salvaterra PM, Lewis J, Hutton M, et al. Tauopathy in *Drosophila*: neurodegeneration without neurofibrillary tangles. *Science* 2001; 293:711–4.
- LaFerla FM, Green KN, Oddo S. Intracellular amyloid-beta in Alzheimer's disease. *Nat Rev Neurosci* 2007; 8:499–509.
- Esselens C, Oorschot V, Baert V, Raemackers T, Spittaels K, Serneels L, et al. Presenilin 1 mediates the turnover of telencephalin in hippocampal neurons via an autophagic degradative pathway. *J Cell Biol* 2004; 166:1041–54.
- Zhang H, Sun S, Herreman A, De Strooper B, Bezprozvanny I. Role of presenilins in neuronal calcium homeostasis. *J Neurosci* 30:8566–80.
- Martin-Morris LE, White K. The *Drosophila* transcript encoded by the beta-amyloid protein precursor-like gene is restricted to the nervous system. *Development* 1990; 110:185–95.
- Leyssen M, Ayaz D, Hebert SS, Reeve S, De Strooper B, Hassan BA. Amyloid precursor protein promotes post-developmental neurite arborization in the *Drosophila* brain. *EMBO J* 2005; 24:2944–55.
- Slunt HH, Thinakaran G, Von Koch C, Lo AC, Tanzi RE, Sisodia SS. Expression of a ubiquitous, cross-reactive homologue of the mouse beta-amyloid precursor protein (APP). *J Biol Chem* 1994; 269:2637–44.
- Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, et al. A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature* 2007; 448:151–6.
- Williamson WR, Hiesinger PR. Preparation of developing and adult *Drosophila* brains and retinae for live imaging. *J Vis Exp* 2010.